


POLYPHENOL METABOLITE LIBRARY OF STANDARDS (PPMLS)

Cat No. PPMLS



CONVENIENT 96-WELL FORMAT Easy storage, no glass bottles
Plated to allow row multiplexing for efficient processing

HIGH PURITY and STABLE
Supplied as 5 µg dried weight; plenty of material for multiple injections

MLSDiscovery™ SOFTWARE
Data processing, data collection and data reduction tool creates libraries in hours

PPMLS™ (Polyphenol Metabolite Library of Standards) is a collection of high-quality small biochemicals that have a polyphenol structure (i.e., several hydroxyl groups on aromatic rings). Polyphenols are found in many plant foods including fruits, vegetables, cereals, dry legumes, chocolate, oils, spices and some beverages, including red wine. Polyphenols are a class of compounds which include flavonoids, phenolic acids, lignans, and stilbenes. In plants, these secondary metabolites are generally involved in stress defense. Studies of polyphenol compounds have shown evidence of anti-oxidative properties and their consumption may limit the risk of degenerative diseases associated with oxidative stress and inflammation.

These are high purity (>95%) compounds supplied in an economical, ready-to-use format. These standards are most commonly used to provide retention times and spectra for these key metabolic compounds, help optimize mass spectrometry analytical protocols, and qualify and quantify mass spectrometry sensitivity and limit of detection.

PPMLS comes with **MLSDiscovery™**, a software tool to support the extraction, manipulation, and storage of the data generated when using Libraries of authentic metabolite standards produced by IROA Technologies.

Features and Benefits

Compounds

PPMLS contains 80 unique small molecule metabolites covering key metabolic pathways, including the following classes of compounds:

- Phenols
- Phenolic terpenes
- Phenylpropanes
- Flavonoids, Isoflavonoids. Flavanones and Methoxyflavonols
- Cinnamic acids and derivatives
- Quinic acids and derivatives
- Stilbenes
- Lignans

Convenient

- High purity metabolites, pre-weighed, supplied dried, 5µg per compound
- Ideal for mass spectrometry metabolomics applications

Formatted

PPMLS contains 80 unique small molecule metabolites.

- Arrayed in (1) one 96-well plate
 - Greiner MASTERBLOCK® #780215, polypropylene deep-well (total volume per well = 1.2mL) in combination with Agilent 24214-001 Peelable PlateLoc Heatseals
- Suitable for manual and automated workflow

Software

MLSDiscovery software package is distributed with and is tailored to work with AAPMLS to help build a physical mass spectral library using the analytical conditions that are normally employed in the user laboratory. MLSDiscovery collects spectra, retention time, mass and relative intensity information for compounds, fragments and adducts.

The software supports most MS data files including Agilent and Thermo. The requirements of the program are that:

- 1) The computer should have at least 8 GB of RAM; Windows 7 or higher.
- 2) Data files must be able to be converted to mzXML format.

To facilitate the use of the program a MLSDiscovery User Manual is provided to help you run through the standard workflow.

Plate Map

The library compounds are arranged in (1) one 96-well polypropylene rack with alphanumeric assigned positions. Please refer to plate map excel sheet that comes with the library for compound locations and identifiers.

Occasionally the map plate will change due to the availability of compounds. Although we try to make sure that the compounds of each row have distinct molecular weights and can be multiplexed, users should refer to the plate map before proceeding.

The plate map contains descriptors and represents information gathered from multiple databases and therefore may contain errors. We suggest that the information provided is carefully reviewed. To help build a better database, please report any discrepancies.

The excel spreadsheet plate map includes identifiers as follows:

Plate number - total number of plates is one (1).

Plate row letter - rows are marked A-H.

Plate column number - columns are numbered 1-12.

Primary compound name – from HMDB, KEGG or PubChem where available.

SMILES – from https://pubchem.ncbi.nlm.nih.gov/search/help_search.html#Smiles

SMILES -- **Simplified Molecular Input Line Entry System**, a chemical structure *line notation* (a typographical method using printable characters) for entering and representing molecules. SMILES strings can be imported or exported from many molecular editors.

Inchikey – a 27-character long identifier based on The IUPAC International Chemical Identifier (InChI™).

Molecular formula – formula of neutral form without salts or water (except in cases where compound has an innate positive charge).

KEGG ID or ChemSpider ID (CSID) – KEGG number where available, otherwise CSID provided.

CAS ID – a unique numerical identifier assigned by Chemical Abstracts Service, a division of the American Chemical Society, to every chemical substance described in the open scientific literature.

HMDB/YMDB ID – Human Metabolome Data Base or Yeast Metabolome Data Base ID number. Either the chemical name or the CAS number was used to search for the HMDB/YMDB ID entry.

Monoisotopic mass

METLIN ID – Scripps Center for Metabolomics and Mass Spectrometry; HMDB, KEGG or CAS number used for METLIN lookup.

PC CID – PubChem Compound Database ID; HMDB, KEGG or CAS number used for PC CID lookup.

PC SID – PubChem Substance Database ID; PC CID, KEGG, Vendor ID, or CAS number used for PC SID lookup.

CHEBI – Chemical Entities of Biological Interest (ChEBI); HMDB, KEGG or CAS number used for ChEBI lookup.

Supplier compound name

Supplier URL

Supplier Cat. No.

Note: Although some of the compounds as seen in the plate wells will be colorless, many dried compounds naturally exhibit color.

Preparation Instructions

Spin down the contents of each 96-well plate in a 96-well plate holder centrifuge, 3000rpm, for 2-3 minutes, to ensure any loose material is deposited at the bottom of each well.

PPMLS compounds are conveniently provided at 5 µg per well, enough for multiple injections. The compounds can either be used as standards and injected individually or mixed in such a way that the entire library may be examined with reasonable efficiency. The compounds in each row *may* all have unique masses; mixing compounds by row *may* allow multiple compounds to be analyzed per injection.

The following are suggestions and dependent on user chromatography and instrumentation.

- 1) Individual injections. As standards, each well represents a single compound; the entire library may be examined in great detail in 80 injections for each of the unique compounds. (Volumes of 250 µL may be considered).
- 2) Simple multiplex injections. If each row is pooled, then the entire collection may be analyzed in 7 injections of simple mixtures. Keep the well volume to 100 µl or less to prevent loss due to dilution and take 5-10 µl of each well for the pooled sample, then inject 2, 4, or 6 µl of the pooled material as needed.

Note: Be sure to check the individual masses across plate rows to ensure these compounds can be separated with the chromatographic system employed.

Solubilization and compound preparation procedure

The following are suggestions and dependent on user chromatography and instrumentation.

- 1) Use high-quality solvents. Compounds in Plate 1 Rows A1-D12 can be solubilized using Ethanol; compounds in Rows E are Water soluble; and compounds in rows F and G are Methanol soluble. Pipet liquid up and down in the well 2-3 times to facilitate solubilization.

- 2) Pool compounds as desired for multiplexing. Again, be sure to check the compound masses you wish to multiplex on the plate map to ensure you can adequately separate the compounds using your chromatographic system prior to pooling.

Precautions and Disclaimer

The PPMLS product is for laboratory research use only. Wear safety glasses and handle with gloves. Avoid contact with skin and eyes. Please consult the Safety Data Sheet for safe handling practices and hazards information.

Storage/Stability

Store plate at -20° C. Once diluted the plate should be resealed and kept at -20° C or -80° C for long-term storage and protected from moisture and light. Avoid repeated freeze/thaw cycles.

Metabolite Libraries Available from IROA

MS Metabolite Library of Standards (MSMLS) - Over 600 unique compounds arrayed in seven (7) 96-well plates that span a broad range of primary metabolism; 5 µg per well.

Large Scale Metabolite Library of Standards (LSMLS) - Over 500 unique compounds arrayed in seven (7) 96-well plates that span a broad range of primary metabolism; 1 mg per well.

Bile Acid Carnitine Sterol Metabolite Library of Standards (BACSMLS) - 96 bile acid, carnitine and sterol metabolites covering key metabolic pathways; 5 µg per well.

Fatty Acid Metabolite Library of Standards (FAMLS) - 96 unique small molecule fatty acid metabolites covering key metabolic pathways; 5 µg per well.

Organic Acid Metabolite Library of Standards (OAMLS) - 96 unique small molecule organic acid metabolites covering key metabolic pathways; 5 µg per well.

Amino Acid/Peptide Metabolite Library of Standards (AAPMLS) – 96 unique metabolites including acetylated, methylated and hydroxy amino acids and dipeptides which are building blocks of proteins in many prokaryotic and eukaryotic organisms; 5 µg per well.

Microbiome Metabolite Library of Standards (GUTMLS) – 185 unique small biochemicals that the gut microbiome produces and interacts with including bacterial, dietary and host xenobiotic metabolites; 5 µg per well.

Phytochemical Metabolite Library of Standards (PHYTOMLS) - 364 unique primary and secondary plant metabolites obtained from consuming diets containing fruits, vegetables, whole grains, legumes, nuts, and plant-based beverages; 5 µg per well.

Polyphenol Metabolite Library of Standards (PPMLS) – 80 unique high-quality small biochemicals that have a polyphenol structure (i.e., several hydroxyl groups on aromatic rings). Polyphenols are found in many plant foods including fruits, vegetables, cereals, dry legumes, chocolate, oils, spices, and some beverages, including red wine.

Legal Information

BACSMLS, AAPMLS, OAMLS, FAMLS, GUTMLS, PHYTOMLS, PPMLS, MSMLS, LSMLS and MLSDiscovery are trademarks of IROA Technologies LLC.

MasterBlock is a trademark of Greiner Bio-One GmbH.

Acknowledgements

We gratefully acknowledge usage of the following websites and databases for their publicly accessible information.

Database	Website
The Human Metabolome Database (HMDB), v 2.5 [1-3]	http://www.hmdb.ca/
The Yeast Metabolome Database [4]	http://www.ymdb.ca/
Chemical Entities of Biological Interest (ChEBI) [5]	https://www.ebi.ac.uk/chebi/
Chemical Abstracts Service (CAS) REGISTRY Database [6]	https://www.cas.org/
Kyoto Encyclopedia of Genes and Genomes (KEGG) [7]	http://www.genome.jp/kegg/
The METLIN Metabolomics Database [8-9]	http://metlin.scripps.edu/index.php
The PubChem Compound and Substance Database [10]	https://pubchem.ncbi.nlm.nih.gov/

References

- [1] Wishart DS, Tzur D, Knox C, et al., HMDB: the Human Metabolome Database. Nucleic Acids Res. 2007 Jan;35 (Database issue):D521-6. 17202168.
- [2] Wishart DS, Knox C, Guo AC, et al., HMDB: a knowledgebase for the human metabolome. Nucleic Acids Res. 2009 37(Database issue):D603-610. 18953024.
- [3] Wishart DS, Jewison T, Guo AC, Wilson M, Knox C, et al., HMDB 3.0 — The Human Metabolome Database in 2013. Nucleic Acids Res. 2013. Jan 1;41(D1):D801-7. 23161693.

- [4] Jewison T, Neveu V, Lee J, Knox C, Liu P, Mandal R, Murthy RK, Sinelnikov I, Guo AC, Wilson M, Djoumbou Y and Wishart DS. YMDB: The Yeast Metabolome Database. *Nucleic Acids Res.* 2012 Jan; 40(Database Issue): D815-20 PubMed: 22064855.
- [5] Hastings J, de Matos P, Dekker A, Ennis M, Harsha B, Kale N, Muthukrishnan V, Owen G, Turner S, Williams M, Steinbeck C. The ChEBI reference database and ontology for biologically relevant chemistry: enhancements for 2013. *Nucleic Acids Res.* 2013. Jan;41(Database issue):D456-63. doi: 10.1093/nar/gks1146. Epub 2012 Nov 24.
- [6] CAS REGISTRY, Division of the American Chemical Society
- [7] Kanehisa M, Goto S. "KEGG: Kyoto Encyclopedia of Genes and Genomes". *Nucleic Acids Res.* 2000 28 (1): 27-30. doi:10.1093/nar/28.1.27. PMC 102409.PMID 10592173.
- [8] Tautenhahn R, Cho K, Uritboonthai W, Zhu Z, Patti G, Siuzdak G. An accelerated workflow for untargeted metabolomics using the METLIN database. *Nature Biotechnology* 2012 30: 826-828. doi:10.1038/nbt.2348.
- [9] Smith CA, I'Maille G, Want EJ, Qin C, Trauger SA, Brandon TR, Custodio DE, Abagyan R, Siuzdak G. METLIN: a metabolite mass spectral database. *The Drug Monit* 2005 27 (6): 747-51. doi:10.1097/01.ftd.0000179845.53213.39. PMID 16404815.
- [10] Kim S, Thiessen PA, Bolton EE, Chen J, Fu G, Gindulyte A, Han L, He J, He S, Shoemaker BA, Wang J, Yu B, Zhang J, Bryant SH. PubChem Substance and Compound databases. *Nucleic Acids Res.* 2016 Jan 4; 44(D1):D1202-13. Epub 2015 Sep 22 [PubMed PMID: 26400175] doi: 10.1093/nar/gkv951 [Free Full Text at Oxford Journals]

IROA MLS Referenced Peer-Reviewed Papers

Diamantidou D, Sampsonidis I, Liapikos T, Gika H, Theodoridis G. Liquid chromatography-mass spectrometry metabolite library for metabolomics: Evaluating column suitability using a scoring approach, *J. Chromatogr. A.* 2023 Feb 8;1690:463779. doi: 10.1016/j.chroma.2023.463779. Epub 2023 Jan 11. PMID: 36681007

Fraisier-Vannier O, Chervin J, Cabanac G, et al. MS-CleanR: A Feature-Filtering Workflow for Untargeted LC-MS Based Metabolomics. *Anal Chem.* 2020;92(14):9971-9981. doi:10.1021/acs.analchem.0c01594. PMID: 32589017

Misra BB, Olivier M. High Resolution GC-Orbitrap-MS Metabolomics Using Both Electron Ionization and Chemical Ionization for Analysis of Human Plasma [published online ahead of print, 2020 Feb 10]. *J Proteome Res.* 2020;10. doi:10.1021/acs.jproteome.9b00774. PMID: 31978300

Vargas F, Weldon KC, Sikora N, et al. Protocol for Community-created Public MS/MS Reference Spectra Within the Global Natural Products Social Molecular Networking Infrastructure [published online ahead of print, 2020 Jan 13]. *Rapid Commun Mass Spectrom.* 2020; e8725. doi:10.1002/rcm.8725. PMID: 31930757

Nye LC, Williams JP, Munjoma NC, et al. A comparison of collision cross section values obtained via travelling wave ion mobility-mass spectrometry and ultra high performance liquid chromatography-ion

mobility-mass spectrometry: Application to the characterisation of metabolites in rat urine. *J Chromatogr A*. 2019 Sept ;1602:386–396. doi:10.1016/j.chroma.2019.06.056. [PMID: 31285057](#)

Dueñas ME, Larson EA and LEE JY. Toward Mass Spectrometry Imaging in the Metabolomics Scale: Increasing Metabolic Coverage Through Multiple On-Tissue Chemical Modifications. *Front. Plant Sci*. 2019 July 10;860. Doi 10.3389/fpls.2019.00860. [PMID: 31354754](#)

Pezzatti J, González-Ruiz V, Codesido S, Gagnebin Y, Joshi A, Guillarme D, Schappler J, Picard D, Boccard J, Rudaz S. A scoring approach for multi-platform acquisition in metabolomics. *J Chromatogr A*. 2019 May 10;1592:47-54. doi: 10.1016/j.chroma.2019.01.023. Epub 2019 Jan 10. [PMID: 30685186](#)

González-Ruiz V, Schwartz D, Sandström J, Pezzatti J, Jeanneret F, Tonoli D, Boccard J, Monnet-Tschudi F, Sanchez JC, Rudaz S. An Integrative Multi-Omics Workflow to Address Multifactorial Toxicology Experiments. *Metabolites*. 2019 Apr 24;9(4). pii: E79. doi: 10.3390/metabo9040079. [PMID: 31022902](#)

Nichols CM, Dodds JN, Rose BS, et al. Untargeted Molecular Discovery in Primary Metabolism: Collision Cross Section as a Molecular Descriptor in Ion Mobility-Mass Spectrometry. *Analytical Chemistry*. 2018 Dec;90(24):14484-14492. DOI: 10.1021/acs.analchem.8b04322. [PMID: 30449086](#)

Pimentel G, Burton KJ, von Ah U, Bütikofer U, Pralong FP, Vionnet N, Portmann R, Vergères G. Metabolic Footprinting of Fermented Milk Consumption in Serum of Healthy Men. *J Nutr*. 2018 Jun 1;148(6):851-860. doi: 10.1093/jn/nxy053. [PMID: 29788433](#)

Thomason K, Babar MA, Erickson JE, Mulvaney M, Beecher C, MacDonald G. Comparative physiological and metabolomics analysis of wheat (*Triticum aestivum* L.) following post-anthesis heat stress. *PLoS ONE* 2018 June13(6): e0197919. <https://doi.org/10.1371/journal.pone.0197919>.

Nemkov T, Hansen KC, D'Alessandro A. A three-minute method for high-throughput quantitative metabolomics and quantitative tracing experiments of central carbon and nitrogen pathways. *Rapid Commun Mass Spectrom*. 2017 Apr 30;31(8):663-673. doi: 10.1002/rcm.7834. [PMID: 28195377](#)

Depke T, Franke R, Brönstrup M. Clustering of MS² spectra using unsupervised methods to aid the identification of secondary metabolites from *Pseudomonas aeruginosa*. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2017 Dec 15;1071:19-28. doi: 10.1016/j.jchromb.2017.06.002. Epub 2017 Jun 4. [PMID: 28642031](#)

González-Ruiz V, Pezzatti J, Roux A, Stoppini L, Boccard J, Rudaz S. Unravelling the effects of multiple experimental factors in metabolomics, analysis of human neural cells with hydrophilic interaction liquid chromatography hyphenated to high resolution mass spectrometry. *J Chromatogr A*. 2017 Dec 8;1527:53-60. doi: 10.1016/j.chroma.2017.10.055. Epub 2017 Oct 25. [PMID: 29106965](#)

Lu, X., Solmonson, A., Lodi, A. et al. The early metabolomic response of adipose tissue during acute cold exposure in mice. *Sci Rep* 2017 June 7; 3455. doi.org/10.1038/s41598-017-03108-x. [PMID: 27399036](#)

Korte AR, Stopka SA, Morris N, Razunguzwa T, Vertes A. Large-Scale Metabolite Analysis of Standards and Human Serum by Laser Desorption Ionization Mass Spectrometry from Silicon Nanopost Arrays. *Anal Chem*. 2016 Sep 20;88(18):8989-96. doi: 10.1021/acs.analchem.6b01186. Epub 2016 Jul 22. [PMID: 27399036](#)